

Advancing High Speed MIR Detection with Quantum Cascade Detectors


The mid-infrared (MIR) region of the electromagnetic spectrum stands out for its vast potential across numerous applications. It boasts rich rotovibrational spectra of various light molecules (small organic molecules, gases, etc.), making **MIR absorption spectroscopy** an invaluable tool for **label-free detection in diverse fields**. Additionally, MIR wavelengths exhibit low scattering by aerosols, rendering them highly promising for research in free-space communication. Notably, specific regions within the MIR spectrum (around 4 μm and 10 μm) offer low absorption by atmospheric gases, facilitating long-distance free-space communication.

The success of MIR applications is rooted in the availability of the required MIR photonic technologies. As such, quantum cascade detectors (QCDs) emerge as a key technology. These photovoltaic detectors are designed to operate over different spectral regions of the MIR. What sets them apart is their ability to **function at room temperature without a bias voltage**. Furthermore, they are characterized by their **low noise**, which compensates for their relatively lower photoresponse compared to alternative MIR detectors. This characteristic pushes their **specific detectivity above $1 \times 10^9 \text{ cm} \cdot \text{Hz}^{1/2}/\text{W}$** . However, the most interesting feature of QCDs is their **exceptional speed**, theoretically exceeding 100 GHz and often exceeding 20 GHz at a -3 dB threshold.

Hamamatsu Photonics is proud to have released **the world's first commercially available QCD^[1,2,3]**, marking a significant milestone in MIR technology.



Hamamatsu Photonics' quantum cascade photodetector (QCD) P16309-01.



Remarkably, this groundbreaking device is **one of the only commercial QCDs operating at room temperature without necessitating any cooling mechanism**. Its applications include high-speed detection of gases^[4] and high-speed spectroscopy^[5] in the MIR region. For example, QCDs can potentially play a critical role in enabling kinetic studies of chemical reactions, which often occur at sub-nanosecond time scales. This capability enables the development of new chemical processes, impacting various facets of life, from **improving energy yield to reducing emissions and promoting the adoption of eco-friendly chemicals**.

Moreover, the high speed of QCDs paves the way for realizing **free-space communication in the MIR region**. Their small size and hassle-free operation bring them closer to widespread adoption in large-scale applications such as communication. Beyond these applications, many more MIR applications stand to benefit from the impressive performance parameters and simplified packaging of QCDs.

For further information, Hamamatsu's dedicated team of engineers is available to answer your queries at info@hamamatsu.eu.

References

^[1] <https://www.hamamatsu.com/jp/en/news/products-and-technologies/2021/20210928000000.html>

^[2] <https://www.hamamatsu.com/jp/en/product/optical-sensors/infrared-detector/qcd/P16309-01.html>

^[3] <https://doi.org/10.1063/5.0038147>

^[4] <https://doi.org/10.3390/s21175706>

^[5] <https://doi.org/10.1038/s42005-020-00420-3>